

**ALLELOPATHIC EFFECT OF AQUEOUS EXTRACTS OF *Tithonia rotundifolia* P.M.
Blake AND *Murraya koenigii* L. ON THE GROWTH AND BIOCHEMICAL
CONSTITUENTS OF *Capsicum annum* L. AND *Corchorus olitorius* L.**

TIJANI MUSA OYEBAMIJI

2012

**ALLELOPATHIC EFFECT OF AQUEOUS EXTRACTS OF *Tithonia rotundifolia* P.M.
Blake AND *Murraya koenigii* L. ON THE GROWTH AND BIOCHEMICAL
CONSTITUENTS OF *Capsicum annuum* L. AND *Corchorus olitorius* L.**

BY

TIJANI MUSA OYEBAMIJI

B.Sc (Ed.) Biology (Ife)

**A THESIS SUBMITTED TO THE DEPARTMENT OF BOTANY, FACULTY OF
SCIENCE, OBAFEMI AWOLOWO UNIVERSITY, ILE-IFE,**

**IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE AWARD OF THE
DEGREE OF MASTERS OF SCIENCE M.Sc. BOTANY**

2012

OBAFEMI AWOLOWO UNIVERSITY

HEZEKIAH OLUWASANMI LIBRARY

POSTGRADUATE THESIS

AUTHORIZATION TO COPY

Author: TIJANI, Musa Oyebamiji

Title: Allelopathic Effect of Aqueous extracts of *Tithonia rotundifolia* (P.M. Blake) and *Murraya koenigii* L. on the growth and biochemical constituents of *Capsicum annum* L. and *Corchorus olitorius* L.

Degree: M.Sc

Year: 2012

I, Tijani Musa Oyebamiji, hereby authorize the Hezekiah Oluwasanmi Library to copy my thesis, in whole or in part, in response to request from individual researchers or organizations for the purpose of private study or research.

Signature

Date

APPROVAL

This thesis has been approved for the Department of Botany, Obafemi Awolowo
University, Ile-Ife

By

Dr. Olutobi.O. Otusanya
(Supervisor)

Dr. (Mrs.) O. Adedeji
Head
(Department of Botany)

DEDICATION

This thesis is dedicated to my late father, Alhaji Tijani Oyebamiji who did so much to make me what I am today. May his gentle soul rest in peace.

OBAFEMI AWOLowo UNIVERSITY

ACKNOWLEDGEMENT

I want to thank God for His profound help and protection on me to start and accomplish this work successfully. I will forever be grateful unto Him for standing by me throughout my stay in the university.

Words are not enough to express my deep and sincere appreciation to my supervisor, Dr. Olutobi.O. Otusanya for her advice, assistance, time, personal sacrifice and endless patience for the successful completion of this work. May the Almighty Allah continue to be with you and your family, in all your undertakings and grant you all your heart desires excellently. I also wish to credit all the entire staff of Botany Department of Obafemi Awolowo University, Ile-Ife: Prof. A.A. Adelusi, Dr. A.M. Makinde, Dr. S.O. Oke, Dr. H.C. Illoh, Mr. B.E. Ayisire, Dr. A.M.A. Sakpere, Prof. A.O. Isichei, Prof. J.I. Muoghalu, Dr. (Mrs) O. Adedeji, Dr. F.A. Oloyede, Dr. A. Saheed, Dr. Tony Odiwe, Dr. A.E. Folorunso, Dr. O.T. Oladipo and to all non teaching staff of the Department: Mr. Gabriel Ademoriyo, Mrs. A. Adedoyin, Mrs. O. T. Ayeloja, Mr. H.O. Oladejo, Mr. Ibiroke for imparting knowledge in me. It is impossible to acknowledge fully all the sources from which assistance has been received. However, there is a great depth of gratitude to the National Horticultural Research Institute (NIHORT) Ibadan, Oyo State for supplying the seeds of *Corchorus olitorius* and *Capsicum annum* (var. Bawa) used in this work.

I appreciate all my friends and postgraduate students in Department of Botany: Ologundudu A.F., Okunlola Olanrewaju, Ogunwole A.A., Awosika Olakanmi, Hassan Sekinat, Olorungbeja John, Stephen, Olabiran Wole, Oguntoye Lateef, Mudashiru Akofe, Oyewole Oyewumi, Oguntoye Sulaiman, Tijani Alowonle, Banjo. You have all contributed in one way or

the other to what I am today. I pray that God will be with you in His unalloyed love and grant all your heart desires excellently.

I want to say a big thank you to my beloved wife, Mrs Tijani Lateefat Bukola and my young ones: Ibrahim, Islamiyat and Ismaeel for given me the required encouragement, care, co-operation and unconditional understanding. To them, I am sincerely thankful and grateful.

I also wish to thank Mr.O.J. Ilori for his academic contributions. May the Almighty Allah bless you with long life and prosperity.

My acknowledgement will never be complete if I do not remember the contribution of AbdulSalami Bolanle and Tijani Sarafadeen. Indeed, you are wonderful.

TABLE OF CONTENTS

| | Page |
|---|------|
| Approval | ii |
| Dedication | iii |
| Acknowledgement | iv |
| Table of Contents | vi |
| List of Figures | ix |
| List of Plates | xiii |
| Abbreviations | xiv |
| Abstract | xv |
| INTRODUCTION | 1 |
| LITERATURE REVIEW | 4 |
| MATERIALS AND METHODS | 11 |
| 3.1 Plant Materials | 11 |
| 3.1.1 Collection of Plant Materials | 11 |
| 3.2 Raising of Seedlings of <i>Tithonia rotundifolia</i> P.M. Blake | 11 |
| 3.3 Preparation of the fresh shoot aqueous extracts of <i>T. rotundifolia</i> and <i>Murraya koenigii</i> L. | 14 |
| 3.4 Germination Experiment | 14 |
| 3.5 Sowing of Seeds of <i>Capsicum annum</i> L. and <i>Corchorus olitorius</i> L. | 15 |



| | | |
|----------------|--|----|
| 3.6 | Harvest | 15 |
| 3.7 | Measurement of growth Parameters, fresh and Dry weights of the test crops | 16 |
| 3.8 | Determination of Chlorophyll Contents | 17 |
| 3.9 | Determination of Crude Protein Contents | 17 |
| 3.10 | Determination of Ascorbic acid accumulation | 18 |
| 3.11 | Statistical Analysis | 19 |
| RESULTS | | 20 |
| 4.1 | Effect of the application of different concentrations of the fresh shoot aqueous extracts of <i>Murraya koenigii</i> and <i>T. rotundifolia</i> on the germination of <i>C. annuum</i> and <i>C. olitorius</i> seeds | 20 |
| 4.2 | Effect of the application of different concentration of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on the plumule length of the germinating seedlings of the test crops | 22 |
| 4.3 | Change in radicle length of the germinating seedlings of the test crops as affected by the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> | 24 |
| 4.4 | Effect of the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on some growth parameters, fresh and dry matter production of the test crops | 26 |
| 4.5 | Effect of the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on chlorophyll accumulation of the test crops | 51 |
| 4.6 | Effect of the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on the crude protein content in the shoots of the test crops | 60 |

| | | |
|--------------------------------------|--|----|
| 4.7 | Effect of the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on the crude protein content in the fruits of the test crops | 63 |
| 4.8 | Effect of the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on ascorbic acid accumulation in the shoot of the test crops | 65 |
| 4.9 | Effect of the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on ascorbic acid accumulation in the fruits of the test crops | 69 |
| DISCUSSION | | 71 |
| CONCLUSION AND RECOMMENDATION | | 79 |
| REFERENCES | | 81 |
| APPENDICES | | 90 |

LIST OF FIGURES

| FIG. | | Page |
|-------------|--|-------------|
| 1. | The effect of the application of fresh shoot aqueous extracts of <i>Murraya koenigii</i> and <i>Tithonia rotundifolia</i> on the germination of seeds of <i>Capsicum annuum</i> and <i>Corchorus olitorius</i> | 21 |
| 2. | Effects of the application of different concentrations of the fresh shoot Aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on the plumule length of the test crops | 23 |
| 3. | Change in the radicle length of germinating seedlings of the test crops as affected by the application of different concentrations of the fresh shoot aqueous extracts of <i>koenigii</i> and <i>T. rotundifolia</i> . | 25 |
| 4a. | The effect of the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> on the shoot height of <i>C. annuum</i> plants. | 27 |
| 4b. | The change in the shoot height of <i>C. olitorius</i> plants subjected to the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> treatments | 28 |
| 5a | Variation in the root length of <i>C. annuum</i> plants as induced by the application of the fresh shoot aqueous extracts of <i>M. koenigii</i> and <i>T. rotundifolia</i> | 30 |

- 5b Variation in the root length of *C. olitorius* plants as affected by the application of fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 31
- 6a The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the number of leaves on *C. annuum* plants 33
- 6b The variation in the number of leaves on *C. olitorius* plants as affected by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 34
- 7a The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the leaf area of *C. annuum* plants 36
- 7b The variation in the leaf area of *C. olitorius* plants as induced by application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 37
- 8a The change in the leaf area ratio of *C. annuum* plants as induced by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 39
- 8b The change in the leaf area ratio of *C. olitorius* plants as affected by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 40
- 9a The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the shoot fresh weight of *C. annuum* plants 42
- 9b The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the shoot fresh weight of *C. olitorius* plants 43
- 10a Variations in the root fresh weight of *C. annuum* plants as induced by the application



- of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 44
- 10b Variations in the root fresh weight of *C. olerarius* plants as induced by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 45
- 11a Change in the shoot dry weights of *C. annuum* plants as affected by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 47
- 11b Change in the shoot dry weights of *C. olerarius* plants as affected by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 48
- 12a Change in the root dry weights of *C. annuum* plants as affected by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 49
- 12b Change in the root dry weights of *C. olerarius* plants as affected by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 50
- 13a The time-course of chlorophyll a accumulation in the shoots of *C. annuum* plants subjected to treatment with the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 52
- 13b The time-course of chlorophyll a accumulation in the shoot of *C. olerarius* plants as affected by the application of the fresh shoot aqueous extracts of *M. keonigii* and *T. rotundifolia* 53
- 14a The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the chlorophyll b accumulation in the shoot of *C. annuum*



- plants.
55
- 14b The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the chlorophyll b accumulation in the shoot of *C. olitorius* plants
56
- 15a The time-course of total chlorophyll accumulation in the shoot of *C. annuum* plants as affected by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 58
- 15b The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the total chlorophyll accumulation in the shoot of *C. olitorius* 59
- 16a Variation in the crude protein accumulation of *C. annuum* plants as induced by the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* 61
- 16b The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the crude protein accumulation in the shoot of *C. olitorius* plants.
62
- 17 The effect of the application of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the crude protein contents in the fruits of the test crops
64
- 18a The change in the ascorbic acid contents of *C. annuum* plants as induced by

- the application of the fresh shoot aqueous extracts of *M. koenigii* and
T. rotundifolia 66
- 18b The time-course of ascorbic acid accumulation in the shoot of *C. olitorius* plants
as affected by the application of the fresh shoot aqueous extracts of *M. koenigii*
and *T. rotundifolia* 68
- 19 Variation in the ascorbic acid contents of the fruits of the test crops as induced by
the application of the fresh shoot aqueous extracts of *M. koenigii* and
T. rotundifolia 70

LIST OF PLATES

Plate

- | | | |
|---|--|----|
| 1 | Seedlings of <i>Tithonia rotundifolia</i> in the nursery | 12 |
| 2 | Seedlings of <i>Murraya koenigii</i> in the nursery | 13 |

OBAFEMI AWOLOWO UNIVERSITY

ABBREVIATIONS

The following abbreviations were adopted in the investigation of germination, growth parameters, chlorophyll, ascorbic acid accumulation and crude protein content determination

CTR: Control supplied with tap water

MPG: Mean Percentage Germination of the test crops

FSEM: Test crops to which fresh shoot aqueous extract of *M. koenigii* was applied

FSET: Test crops to which fresh shoot aqueous extract of *T. rotundifolia* was applied

LAR: Leaf Area Ratio of the test crops

ABSTRACT

The study was conducted to investigate the allelopathic effects of fresh shoot aqueous extracts of *Murraya koenigii* L. and *Tithonia rotundifolia* P.M. Blake on the growth and biochemical constituents of *Capsicum annum* L. and *Corchorus olitorius* L. plants.

Germination experiment was carried out by raising the seedlings of the two target crops in Petri-dishes lined with Whatman No 1 filter paper and moistened with 10 ml of different concentrations (50% and 100% representing half and full strength) of the aqueous extracts of *M. koenigii* and *T. rotundifolia*. For the growth parameters (shoot height, root length, number of leaves, leaf area, leaf area ratio), yield parameters (fresh shoot and root weight, dry shoot and root weight) and quality parameters (chlorophyll, ascorbic acid and crude protein) analyses, potted plants were separated into the control and two other regimes namely: fresh shoot aqueous extract of *T. rotundifolia* (FSET) and fresh shoot aqueous extract of *M. koenigii* (FSEM). The pots were arranged in a completely randomized design. The control plants were supplied with 600 ml of tap water while the extract-treated plants were supplied with 600 ml of the appropriate aqueous extracts daily. Harvesting started at two weeks and continued thereafter on a weekly basis for six weeks. Chlorophyll accumulation, ascorbic acid, percentage nitrogen and crude protein content were determined using standard methods. The data were analyzed using Analysis of Variance (ANOVA) and Least Significance Difference (LSD $p < 0.05$).

The extracts significantly inhibited the germination of the seeds and the plumule and radicle lengths of both target crops. This effect was extract concentration dependent (100% > 50% > Control). The applied extracts of *M. koenigii* plants significantly promoted virtually all the growth parameters such as shoot height, number of leaves, leaf area, leaf area ratio, fresh

shoot weight, fresh root weight, dry shoot weight, dry root weight, chlorophyll a, chlorophyll b, total chlorophyll, ascorbic acid and protein accumulation of the two target crops. The aqueous extract of *T. rotundifolia* enhanced only the shoot height, leaf area, root fresh weight, chlorophyll a, chlorophyll b, total chlorophyll, ascorbic acid and protein accumulation in the shoot of *C. annuum*. The root length was however, significantly inhibited by both aqueous extracts. In the case of the potted plants, the effects of the extracts on the various parameters studied followed the order: FSEM > FSET > CONTROL and was target species dependent.

The results presented in this work showed that FSEM had more pronounced stimulatory effects on the studied parameters than the FSET. It was evident that while the level of the allelochemicals in the extracts of the two donor plants was phytotoxic to and inhibited the germination and growth of the juvenile seedlings in the Petri-dishes, they, however, had a stimulatory effect on the growth of the matured potted plants as well as on the accumulation of the biochemical constituents studied.

CHAPTER ONE

INTRODUCTION

The genus *Tithonia* belongs to the family Asteraceae. There are about 11 species of the genus worldwide out of which two are in Nigeria (Muoghalu and Chuba, 2005). *Tithonia rotundifolia* P.M. Blake and *Tithonia diversifolia* (Hemsl.) A. Gray are invasive annual weeds known to be native to Mexico and Central America (Akobundu and Agyakwa, 1987). According to Ayeni *et al.* (1997), they were initially introduced to improve soil fertility and crop yields in Nigeria. However, the weeds are now observed to grow aggressively along road paths, abandoned farmlands and hedges all over Nigeria (Akobundu and Agyakwa, 1987). *Tithonia* species have been reported to contain some allelochemicals and therefore suggested as being capable of posing a serious threat of phytotoxicity to agricultural crops (Tongma *et al.*, 1998).

Olabode *et al.* (2009) found that *T. diversifolia* contains phytochemical constituents such as alkaloids, flavonoids, tannins and saponins. Recent phytochemical screening by Otusanya and Ilori (2012) revealed that the methanolic and water extracts of *T. rotundifolia* contained glycosides, tannins, flavonoids, saponins, phenols, terpenoids and alkaloids. Ayeni *et al.* (1997) stated that allelochemicals that are toxic may inhibit shoot/root growth, nutrient uptake, or may attack a naturally occurring symbiotic relationship, thereby destroying the plants usable source of nutrients. According to the same authors, the consequent effects may be inhibited or retarded germination rate, reduced radicle/root or plumule/shoot extension, lack of root hairs, swelling or necrosis of root tips, curling of the root axis, increased number of seminal roots, discolouration, reduced dry weights accumulation and lowered reproductive capacity. Imeokpara and Okusanya

(1994) observed that most farmers find it difficult to manage the infestation of these weeds in most crop fields particularly in rice and maize fields.

Murraya koenigii L. commonly called 'curry leaf' belongs to the family Rutaceae. It is an aromatic, deciduous shrub which is native to India (Satyavati *et al.*, 1987). The plant is used as a spice for its characteristic flavor and aroma and as a flavouring agent in curries and chutneys (Gopalan *et al.*, 1984). In India, it is used both in conventional and traditional medicine to treat various ailments (Arivoli and Tennyson, 2011). Phytochemical constituents such as alkaloids, volatile oils, xanthoxin and carotenes are present in *M. koenigii* (Chakraborty, 1970; Bordner *et al.*, 1972). It was recently observed that in areas where *M. koenigii* grows, the growth of other plants was hampered, hence, it was suspected to possess allelopathic attributes.

The two test crops in this study are; *Corchorus olitorius* L. and *Capsicum annum* L. The genus *Corchorus* is a member of the family Tiliaceae, native to the tropical and sub-tropical regions of the world (Nath, 1976). *C. olitorius* commonly known as wild okra is a tall herbaceous annual vegetable whose nutritious leaves and fruits are widely consumed among rural communities in most parts of Africa (Velempini *et al.*, 2003). In West Africa, it is commonly cultivated and popularly used for soup among people of all classes especially in Nigeria (Oyedele *et al.*, 2006). According to Zakaria *et al.* (2006), wild okra is used in folklore medicine in the treatment of gonorrhoea, chronic cystitis, pain, fever and tumour.

The genus *Capsicum*, as presently perceived, include at least 25 species of vegetables, four of which have been domesticated. *Capsicum annum* L. is the best known domesticated species in the world (Esbaugh, 1993). Navarro *et al.* (2006) stated that *C. annum* originated from northern Latin America and has become an important agricultural crop, not only because of its economic importance, but also for the nutritional value of its fruits. Howard *et al.* (2000)

observed that this vegetable is an excellent source of natural colours and antioxidant compounds. Hence, the intake of these compounds in food is an important health-protecting factor. According to Bramley (2000), *C. annuum* has been observed to be beneficial for prevention of widespread human diseases, including cancer and cardiovascular diseases. It is also known to contain vitamin C, an important compound of pepper fruits which chelates heavy metal ions (Namiki, 1990), reacts with singlet oxygen and other free radicals and suppresses peroxidation (Bielski *et al.*, 1995; Harris, 1996), thereby reducing the risk of arteriosclerosis, cardiovascular diseases and some forms of cancer.

Therefore, the specific objectives of this study were to;

- a) Investigate the effects of fresh shoot aqueous extracts of *M. koenigii* (FSEM) and *T. rotundifolia* (FSET) on the germination and some growth parameters such as the shoot height, number of leaves, leaf area, leaf area ratio and yield parameters such as fresh shoot and root weights, dry shoot and root weights etc. of *C. annuum* and *C. olerius*, and
- b) Investigate the effects of the fresh shoot aqueous extracts of *M. koenigii* and *T. rotundifolia* on the chlorophyll, crude proteins and ascorbic acid accumulation in the shoots and fruits of *C. annuum* and *C. olerius*

The study is expected firstly, to contribute to the understanding of the importance of allelopathy in weed-crop relations and secondly, to determine the nature of interference of allelochemicals in *M. koenigii* and *T. rotundifolia* on the germination, growth and accumulation of some important chemical metabolites in *C. annuum* and *C. olerius*.

OBAFEMI AWOLOWO UNIVERSITY